

Accepted by Astrophysical Journal Letters, 26 June 2002

Angular momentum extraction by gravity waves in the Sun

Suzanne Talon

Département de Physique, Université de Montréal, Montréal PQ H3C 3J7

CERCA, 5160, boul. Décarie, suite 400, Montréal PQ H3X 2H9

Pawan Kumar

Dept. of Astronomy, University of Texas, Austin, TX 78713

Jean-Paul Zahn

LUTH, Observatoire de Paris, 92195 Meudon, France

Abstract

We review the behavior of the oscillating shear layer produced by gravity waves below the surface convection zone of the Sun. We show that, under asymmetric filtering produced by this layer, gravity waves of low spherical order, which are stochastically excited at the base of the convection zone of late type stars, can extract angular momentum from their radiative interior. The time-scale for this momentum extraction in a Sun-like star is of the order of 10^7 years. The process is particularly efficient in the central region, and it could produce there a slowly rotating core.

Subject headings: hydrodynamics — stars: late-type — stars: rotation — Sun: interior — waves

1. Introduction

Angular momentum transport by gravity waves has received much attention recently. While it has first been considered as a key mechanism in the tidal interaction of binary systems (Zahn 1975; Goldreich & Nicholson 1989), it has since been invoked also as an efficient process of momentum redistribution in single stars (Schatzman 1993; Kumar & Quataert 1997; Zahn, Talon & Matias 1997). The mechanism proposed was similar to that acting in binary stars, with synchronization proceeding gradually inwards. However, as pointed out by Gough & McIntyre (1998) and Ringot (1998), the treatment of angular momentum extraction as presented by these first studies was incorrect, and it became clear that the actual properties of wave transport were far more complex than originally anticipated.




Fig. 1.— Angular momentum luminosity integrated over $0.1\mu\text{Hz}$ as a function of order ℓ and frequency.

Wave properties have been examined further by Barnes, McGregor & Charbonneau (1998) who showed that magnetic fields transform pure gravity waves into gravito-Alfvén waves, thus modifying their damping behavior. They showed that strong magnetic field may prevent certain waves from propagating.

Kumar, Talon & Zahn (1999, hereafter KTZ) then demonstrated how gravity waves in interaction with shear turbulence may lead to the formation of an oscillating shear layer just below the surface convection zone, analogous to the well studied quasi-biennial oscillation of atmospheric sciences (see *e.g.* Shepherd 2000 for details on the QBO and other wave-driven oscillations).

Kim & McGregor (2001) further studied that oscillation in a simplified two waves model. They showed that, with only a prograde and a retrograde wave, and with the velocity fixed both at the top and at the bottom of the solar tachocline, the time-dependent behavior of the rotation profile goes from being periodic, to quasi-periodic and to chaotic as the viscosity in the shear region is slowly decreased. Their goal was to seek an explanation of a 1.3 yr variation, which Howe et al. (2000) claim to have detected in the tachocline.

Here we want to reexamine the effect of waves in the deep solar interior. Indeed, while the high degree waves are damped very close to the base of the convection zone, thus leading to rapid oscillations that could explain rotational velocity variations in the tachocline (Kim & McGregor 2001) or be related to the solar cycle (KTZ), the low degree waves that are also excited stochastically by the convective motions, although with a somewhat lower efficiency (see Kumar & Quataert 1997 and Zahn, Talon & Matias 1997) will have, over a much longer period, an effect on the deep interior.

In this letter, we wish to demonstrate how these waves, in conjunction with shear turbulence, indeed achieve momentum extraction in the deep interior. In §2, we explain heuristically how that process takes places and in §3, we show the results of a long numerical simulation applied to the solar case.




Fig. 2.— Oscillating shear layer below the surface convection zone. The dotted line shows the initial rotation profile. With the surface rotating slower than the core, a prograde shear layer is initially formed, followed by a retrograde one. When the shear becomes too intense, turbulent viscosity acts to merge the prograde layer with the convection zone, leaving behind the retrograde layer. A new prograde layer forms behind, and migrates towards the convection zone when the retrograde layer is absorbed. The cycle then resumes.

2. Shear layer and momentum extraction

The formation of a shear layer is easy to describe, using a simple two waves model. When both a prograde and a retrograde wave travel in an accelerating region (*i.e.* where Ω increases inwards with depth), the prograde wave frequency diminishes, enhancing its damping; the opposite occurs for the retrograde wave (see KTZ for details). This creates a double peaked shear layer. The gradient in that layer rises until it becomes steep enough so that turbulence will start playing a role and will effectively merge the shear layer with the adjacent convection zone. The retrograde layer¹ will then slowly rise toward the convection zone, and a prograde layer will form further inside the star. The retrograde layer is eventually also absorbed by the convection zone, and the cycle goes on.

The oscillating shear layer will play an important role in filtering the (low degree) waves that will penetrate, and thus deposit their momentum in the deep interior. Indeed, when the prograde peak is larger, it is the retrograde waves that will penetrate and deposit their negative momentum in the interior, and vice versa if the retrograde peak dominates. If there is no initial differential rotation in the interior, the shear layer will oscillate symmetrically, and on average no net momentum flux will occur. However, if the surface convection zone is slowed down, as in the case of the Sun, there exists a bias in the formation of the shear layer, which will favor the prograde peak (the interior

¹The retrograde layer is the one that rotates more slowly than the convection zone, while the prograde layer rotates more rapidly.




Fig. 3.— Global evolution of the rotation profile over large time periods. Efficient momentum extraction is visible in a characteristic time-scale of 10^5 years.

then rotating faster than the surface), thus letting a larger amount of negative momentum being transported to the deep interior. The positive momentum deposited in excess in the shear layer will then be transferred back in the surface convection zone by turbulence. The net effect will thus be to extract momentum from the radiation zone.

3. A numerical simulation

While it is clear that the filtering bias introduced by the slope of the rotation profile below the convection zone leads to momentum extraction, we further need to know where exactly this extraction takes place, and how efficient it is. To answer this question, we performed a numerical simulation covering a very long time period. There are two key ingredients in such a simulation. Firstly, one needs to include a whole variety of radial orders to describe both the waves responsible for the formation and oscillation of the shear layer and the deeply penetrating waves. Furthermore, retaining only two wave orders has the effect of depositing momentum too locally, compared to having a more complete spectrum. We thus included spherical orders $\ell = 1, 5, 9, \dots, 57$, to capture a wide variety of damping behaviors. We also considered a small frequency range $\omega = 0.6$ to $1.0 \mu\text{Hz}$. Higher frequency waves will have little impact since the power spectrum we are using in these simulations falls off as $\omega^{-4.5}$ (*cf.* Eq. 5 and KTZ), while lower frequency waves will be damped very close to the convection zone (*cf.* Eq. 3), not undergoing differential damping. The second key ingredient is to follow both short time-scales (~ 10 years), characteristic of the shear layer oscillation, and long time-scales ($\sim 10^7$ years), characteristic of the spin-down rate. This

Table 1: Damping length d_ω (scaled by the solar radius) at 0.6 μHz

ℓ	m	d_ω^a	d_ω^b
1	1	0.616	0.461
	-1		0.661
5	5	0.084	0.036
	-5		0.499
9	9	0.033	0.017
	-9		0.378
13	13	0.018	0.010
	-13		0.256
17	17	0.012	0.007
	-17		0.061
21	21	0.008	0.005
	-21		0.024

^aIn solid body rotation

^bWith the initial rotation profile (*cf.* Figures 2 & 3)

combinaison requires the use of a conservative numerical scheme.

We thus calculated the evolution of angular momentum in the radiative interior by the combined effect of turbulence and gravity waves:

$$\rho \frac{\partial}{\partial t} [r^2 \Omega] = \frac{1}{r^2} \frac{\partial}{\partial r} \left[\rho \nu_t r^4 \frac{\partial \Omega}{\partial r} \right] - \frac{3}{8\pi} \frac{1}{r^2} \frac{\partial}{\partial r} \mathcal{L}_J(r) \quad (1)$$

where ρ is the density, Ω the angular velocity and ν_t the turbulent viscosity (*cf.* Talon 1997). The local momentum luminosity $\mathcal{L}_J(r)$ is integrated over the whole wave spectrum, each wave being damped by radiative processes and turbulent viscosity (see Eq. 7)

$$\mathcal{L}_J(r) = \sum_{\sigma, \ell, m} \mathcal{L}_{J\ell, m}(r_{zc}) \exp[-\tau(r, \sigma, \ell)] \quad (2)$$

and where the local damping rate takes into account the mean molecular weight stratification

$$\tau(r, \sigma, \ell) = [\ell(\ell + 1)]^{\frac{3}{2}} \int_r^{r_c} (K + \nu_t) \frac{N N_T^2}{\sigma^4} \left(\frac{N^2}{N^2 - \sigma^2} \right)^{\frac{1}{2}} \frac{dr}{r^3} \quad (3)$$

where $N^2 = N_T^2 + N_\mu^2$ is the total Brunt-Väisälä frequency, N_T^2 is its thermal part and N_μ^2 is due to the mean molecular weight stratification (*cf.* Zahn et al. 1997). $\sigma(r, m) = \omega - m[\Omega(r) - \Omega_{zc}]$ is the local (Doppler shifted) frequency, and ω is the frequency in the reference frame of the convection

zone. The damping length d_ω defined by $\tau(d_\omega) = 1$ depends strongly on frequency ω and, in presence of differential rotation, on the signed value of the azimuthal wavenumber m , as shown in Table 1. The evolution equation (1) is solved using a decoupled scheme, in which momentum is first deposited by waves and diffusion is then treated using a perfectly conservative finite element method. After 10^7 time steps of 1 year, the error on global momentum conservation is $\sim 10^{-5}$. This equation is solved throughout the radiation zone. The upper boundary condition expresses the conservation of momentum of the star as a whole, with the convection zone rotating as a solid body. On their way to the core, the waves are reflected when their local frequency σ equals the Brunt-Väisälä frequency

$$\sigma(r, m) = N. \quad (4)$$

They then deposit more momentum as they travel back to the convection zone. Some waves do travel all the way to the core. Since that region contains very little momentum, it is easily spun down (see Figure 3). Waves reaching the center most region (below $r = 0.01 R_\odot$) are reflected back; this innermost core has been taken to rotate as a solid body.

Wave generation is due to Reynolds stresses at the base of the convection zone, and to model its amplitude we followed here the description of Goldreich et al. (1994) (see also KTZ). The energy flux per unit frequency $\mathcal{F}_E$ is then

$$\begin{aligned} \mathcal{F}_E(\ell, \omega) &= \frac{\omega^2}{4\pi} \int dr \frac{\rho^2}{r^2} \left[\left(\frac{\partial \xi_r}{\partial r} \right)^2 + \ell(\ell+1) \left(\frac{\partial \xi_h}{\partial r} \right)^2 \right] \\ &\times \exp \left[-h_\omega^2 \ell(\ell+1)/2r^2 \right] \frac{v^3 L^4}{1 + (\omega\tau_L)^{15/2}}, \end{aligned} \quad (5)$$

where ξ_r and $[\ell(\ell+1)]^{1/2}\xi_h$ are the radial and horizontal displacement wavefunctions which are normalized to unit energy flux just below the convection zone, v is the convective velocity, L is the radial size of an energy bearing turbulent eddy, $\tau_L \approx L/v$ is the characteristic convective time, and h_ω is the radial size of the largest eddy at r with characteristic frequency of ω or greater ($h_\omega = L \min\{1, (2\omega\tau_L)^{-3/2}\}$). The gravity waves are evanescent in the convection zone, the region where they are excited. The above equation was derived under the assumption that the turbulence spectrum is Kolmogorov and it ignores wave excitation resulting from convective overshooting. The momentum flux per unit frequency $\mathcal{F}_J$ is then

$$\mathcal{F}_J(m, \ell, \omega) = \frac{m}{\omega} \mathcal{F}_E(\ell, \omega). \quad (6)$$

In the case of the Sun, there is a whole spectrum of waves, in frequency ω and spherical order ℓ . Their range is given by $\omega \leq N_c$, where N_c is the Brunt-Väisälä frequency at the base of the convection zone², and by $1 \leq \ell \leq \ell_c$ where $\ell_c \sim 60$ is the spherical order characterizing

²This frequency depends on the amount of overshooting that is considered. However, since the power spectrum scales as $\omega^{-4.5}$ (cf. KTZ), its exact value is not crucial.

convection (it corresponds to one pressure scale-height). Assuming that all azimuthal orders m are equally excited, we find that most of the momentum is carried by the low frequency waves. There is significant momentum luminosity in the low order waves that do penetrate deep in the interior (*cf.* Figure 1 and Table 1).

For this first exploration, turbulent viscosity has been taken proportional to the radiative viscosity (ν_{rad}), admittedly a very crude prescription:

$$\nu_t = 10^5 \nu_{\text{rad}} = 10^5 \frac{\chi T}{5c^2 \rho} \quad (7)$$

where χ is the thermal conductivity and c the speed of light. The total energy flux in the considered waves is 6.3×10^{29} erg/s, and the total integrated momentum luminosity is 1.7×10^{36} g cm²/s². This turbulent viscosity operates both on the mean rotation profile and on the waves (*cf.* Eq. 3 and 1).

This simulation is performed in one dimension and we only consider the effect of spherically integrated waves. We thus neglect the latitudinal differential rotation just below the convection zone, as well as the influence of the Coriolis force on wave propagation and wave excitation.

The results are displayed in Figures 2 and 3. The first shows the shear layer oscillation cycle, which occurs here with a period of ~ 300 years. The second shows the evolution of the rotation profile in the deep interior over a much longer period. Momentum extraction is related to the asymmetric filtering by the double shear layer, which is determined by the amount of differential rotation initially present below the convection zone. In our simulation we let Ω increase by $0.15 \mu\text{Hz}$ down to $r = 0.60R_{\odot}$. This profile, applied to the wave spectrum we consider here, yields a differential momentum flux below the shear layer of 2×10^{33} g cm²/s². In 5×10^4 years, this leads to an extraction of 2.5×10^{45} g cm²/s, in good agreement with the numerical simulation which shows an extraction of 4×10^{45} g cm²/s.

As the simulation progresses, the radiation zone decelerates and the convection zone accelerates³; this leads to less differential filtering and therefore momentum extraction gradually slows down. The simulation shows that waves are very efficient in extracting momentum from the deep core, in a time short compared to the main-sequence life time. At later times, the net wave flux decreases, allowing turbulence to smooth the rotation profile. But the turbulent transport will also taper off, due to the build-up of a stabilizing helium gradient. Therefore it is plausible that in the present Sun the central region is rotating slower than the rest of the radiation zone, as suggested by certain helioseismic inversions (Elsworth et al. 1995; Corbard et al. 1997). If this trend is confirmed, it would imply that wave transport is more efficient than magnetic torquing, which has also been invoked for the extraction of angular momentum (*cf.* Mestel & Weiss 1987, Charbonneau & MacGregor 1993).

³In an evolving solar model, the surface convection zone would constantly be spinned down by a magnetic torque and would thus probably never accelerate.

4. Discussion

In this letter, we have shown how gravity waves can extract angular momentum from the radiative core of a solar-like star through differential wave filtering, sufficiently to establish an almost uniform rotation profile in about 10^7 years. In our model, the presence of turbulent transport is essential to yield the proper result; indeed, it is required both to produce the QBO like oscillation, and to smooth the excessive differential rotation produced by waves in the central region.

The evolution of latitudinally averaged angular momentum in the Sun would then proceed as follows. The magnetic torque applied to the convection zone tends to slow it down, establishing a negative gradient of angular velocity, which induces the bias between prograde and retrograde waves. Due to the combined action of wave transport and of turbulent viscosity, the velocity gradient tends to diminish. In our simulation it disappears altogether, but in the real Sun, it would be maintained at some level due to the competition between the extraction of angular momentum from the radiative interior, and its loss by the solar wind: the faster the spin-down, the steeper the gradient, the stronger the extraction. If the extraction of angular momentum by gravity waves is as efficient as we estimate, it will adjust itself such as to provide just the flux of angular momentum which is lost by the wind, with the interior profile of angular velocity being rather flat, except near the center and at the top of the radiation zone.

The presence of a large toroidal magnetic at the base of the convection zone could somewhat lengthen the time-scales mentioned here. Indeed, the magnetic field required to prevent wave propagation at $0.6 \mu\text{Hz}$ is about $(3 \times 10^5 / \ell)$ G (*cf.* KTZ). If fields of a strength as high as 10^5 G are present in that region (*cf.* Fan et al. 1993, Caligari et al. 1995), high ℓ modes could be prevented from propagating, modifying the evolution of the double shear layer. However, since such large fields occur in localized areas, the global effect would rather be one of decreasing the wave flux and thus, simply increase the time scale estimated here.

This initial work must be pursued in many ways. Firstly, one would like to assess the efficiency of this mechanism in an evolving solar model, with magnetic spin-down, in which meridional circulation would also be taken into account in order to verify the scenario we proposed earlier.

Other improvements should be made on the physical description. The effect of the Coriolis force must be included, which could require to step up to two-dimensional simulations. Finally, a more realistic prescription for the turbulent transport must be implemented, before we can conclude that the solar core is rotating significantly slower than the rest of the radiation zone.

Acknowledgments S. Talon was supported by NSERC of Canada and by the Canada Research Chair in Stellar Astrophysics awarded to Prof. G. Fontaine.

REFERENCES

- Barnes, G., MacGregor, K. B. and Charbonneau, P., 1998, ApJ 498, L169
- Caligari, P., Moreno-Insertis, F. and Schüssler, M. 1995, ApJ 441, 886
- Charbonneau, P. and MacGregor, K. B., 1993, ApJ 417, 762
- Corbard, T., Berthomieu, G., Morel, P., Provost, J., Schou, J. and Tomczyk, S., 1997, A&A 324, 298
- Elsworth, Y., Howe, R., Isaak, G. R., McLeod, C. P., Miller, B. A., New, R., Wheeler, S. J. and Gough, D. O., 1995, Nature 376, 669
- Fan, Y., Fisher, G.H. and DeLuca, E.E. 1993, ApJ 405, 390
- Goldreich, P., Murray, N. and Kumar, P., 1994, ApJ 424, 466
- Goldreich, P. and Nicholson, P. D., 1989, ApJ 342, 1079
- Gough, D. O., and McIntyre, M. E., 1998, Nature 394, 755
- Howe, R., Christensen-Dalsgaard, J., Hill, F., Komm, R., Larsen, R. M., Schou, J., Thompson, M. J., Toomre, J., 2000, Science 287, 2456
- Kim, Eun-jin, and MacGregor, K.B., 2001, ApJ 556, L117
- Kumar, P. and Quataert, E., 1997, ApJ 475, L143
- Kumar, P., Talon, S. and Zahn, J.-P., 1999, ApJ 520, 859 (KTZ)
- Mestel, L. and Weiss, N.O., 1987, MNRAS 226, 123
- Ringot, O., 1998, A&A 335, 89
- Schatzman, E., 1993, A&A 279, 431
- Shepherd, T. G., 2000, J. of Atmos. and Solar-Terrestrial Physics 62 (17-18) 1587
- Talon, S., 1997, Thèse de doctorat, Univ. Paris 7
- Zahn, J.-P., 1975, A&A 41, 329
- Zahn, J.-P., Talon, S. and Matias, J., 1997, A&A 322, 320